


# InertSearch™ for LC

Inertsil® Applications

## Analysis of Nitrofurantoin Metabolite by Pre-column HPLC

Data No. LA608-0696

Nitrofurans	Major metabolite
nitrofurantoin	1-aminohydantoin(AHD)
furazolidone	3-amino-2-oxazolidinone (AOZ)
nitrofurazone	Semicarbazide(SEM)
furaltadone	3-amino-5-methylmorpholino-2-oxazolidinone (AMOZ)



### Conditions

**System** : Symbiosis Pharma system (LC mode)  
API 2000™

**Column** : Inertsil ODS-SP (3 μm, 50 x 2.1 mm I.D.)

**Column Cat. No.** : 5020-02812

**Eluent** : A) CH<sub>3</sub>OH  
B) 5 mM CH<sub>3</sub>COONH<sub>4</sub>  
A/B = 30/70 - 5 min - 90/10 - 5 min - 90/10, v/v

**Flow Rate** : 0.2 mL/min

**Col. Temp.** : 40 °C

**Detection** : LC/MS/MS (API 2000™: ESI, Positive, MRM)

**Injection Vol.** : 10 μL

**Analyte** : Nitrofran metabolites

### Analyte:

		Q1	Q3
1.	2-NBA-AHD (10 μg/L)	249.0	133.8
2.	2-NBA-AOZ (10 μg/L)	236.2	134.2
3.	2-NBA-SEM (10 μg/L)	209.1	166.1
4.	2-NBA-AMOZ (10 μg/L)	335.3	291.2

2-NBA : 2-nitrobenzaldehyde