

MORRIS TEST-KIT-GUIDE

INSTRUCTIONS

- POWDERIZE YOUR SAMPLE AND PLACE IT ON A WHITE CERAMIC PLATE *DO NOT USE A TEST TUBE*
- 2. DRIP ONE DROP FROM BOTTLE "A" ON THE SAMPLE
- DRIP ONE DROP FROM BOTTLE "B" ON THE SAMPLE
- 4. STIR THE MIXTURE WITH A TOOTHPICK OR THE SHARP POINT OF A KNIFE FOR A FULL 30 SECONDS
- COMPARE COLOR WITH REACTIONS IN THIS BOOK

SAMPLE SIZE

POWDER: USE 5 MILLIGRAMS - ABOUT THE SIZE OF A DROP OF WATER
PILL: SCRAPE OFF A SMALL AMOUNT - AROUND 1/20TH OF A PILL

WARNING! THE TESTING LIQUID IS HAZARDOUS

DO NOT GET IT ON YOUR SKIN OR IN YOUR EYES - DO NOT INGEST JSE IN A WELL VENTILATED AREA AND WEAR GLOVES & EYE PROTECTION IF POSSIBLE. IF THE BOTTLE BECOMES BLOCKED, DO NOT ATTEMPT TO CLEAR BY SQUEEZING. INSTEAD CLEAR THE BLOCKAGE BY SHAKING VIGOROUSLY OR INSERTING A PAPERCLIP! SIMILAR OBJECT.

KEEP OUT OF HEAT/SUNLIGHT

DO NOT REMOVE ABSORBENT POWDER FROM CONTAINER

INSTRUCTIONS

THIS TEST IS DIFFERENT FROM EVERY OTHER KIT PRODUCED BY THE BUNK POLICE - TWO STEPS ARE CRUCIAL TO ACHIEVE THE DESIRED RESULT - 1. CRUSH / POWDERIZE YOUR SAMPLE COMPLETELY BEFORE TESTING 2. STIR THE SAMPLE FOR A FULL 30 SECONDS BEFORE READING THE RESULT.

WARNING


WHEN YOU FIRST ADD THE DROP FROM BOTTLE "A" - BLUE SPECS WILL APPEAR. IGNORE THIS COLOR AND WAIT UNTIL YOU 1. HAVE ADDED THE DROP FROM BOTTLE "B" 2. HAVE FINISHED STIRRING FOR A FULL 30 SECONDS 3. THE COLOR HAS STOPPED CHANGING - BEFORE READING THE REACTION.




COLOR AFTER ADDING DROP "A" - IGNORE

MORRIS





COCAINE




DCK & ZF-DCK





NO REACTION


THE FOLLOWING SUBSTANCES DO NOT REACT WITH THE MORRIS TEST KIT - MEANING THE COLOR THAT IS PRESENTED WILL BE GREEN AS SEEN ABOVE - THIS LIST IS NOT COMPREHENSIVE. RESEARCH IS STILL ONGOING.


PCP · Z-OXO-PCE · 3-MeO-PCP 3-CI-PCP · 3-MeO-PCE · O-PCE MXPr · MXE · EPE · DPD























SAFETY & STORAGE

WARNING! THE TESTING LIQUID IS A STRONG ACID

AVOID SKIN AND EYE CONTACT - DO NOT INGEST

CONDUCT TESTS IN A WELL VENTILATED AREA

USE GLOVES IF POSSIBLE & WASH HANDS AFTER USE

IF THE BOTTLE BECOMES BLOCKED, DO NOT ATTEMPT TO CLEAR IT BY SQUEEZING. INSTEAD CLEAR THE BLOCKAGE BY SHAKING THE BOTTLE VIGOROUSLY OR INSERTING A PAPERCLIP OR SIMILAR OBJECT IF NECESSARY.

KEEP REFRIGERATED - FROZEN IF POSSIBLE (IF USED OUTDOORS KEEP SHADED OR IN A COOLER)

SHELF LIFE DEPENDS ON STORAGE CONDITIONS IF KEPT REFRIGERATED YOUR KIT CAN LAST SIX MONTHS TO A YEAR (WE DO NOT RECOMMEND USING KITS OVER ONE YEAR OLD)

TIPS & TRICKS

- 1. USE MORE THAN ONE TYPE OF KIT FOR INCREASED ACCURACY
- 2. MAKE SURE TO CONDUCT TESTS IN A WELL LIT AREA
- CRUSH YOUR SAMPLE INTO A POWDER BEFORE TESTING
- **4.** THE FIRST 1 3 SECONDS OF THE REACTION ARE THE MOST IMPORTANT PAY CLOSE ATTENTION DURING THIS TIME OR RECORD THE REACTION ON YOUR PHONE SO YOU CAN REPLAY AND ANALYZE IT AS MANY TIMES AS YOU LIKE.
- 5. THERE ARE MANY SUBSTANCES THAT DO NOT CHANGE COLOR OR REACT IN ANY WAY, SOME OF OUR KITS TAKE ADVANTAGE OF THIS, ALLOWING YOU TO SEE THE REACTIONS OF UNWANTED SUBSTANCES (ADULTERANTS) THAT MIGHT BE IN YOUR SAMPLE ALONG WITH THE DESIRED SUBSTANCE.
- 6. OUR REACTION CHARTS ARE MADE USING TIME-LAPSE PHOTOS OF VERIFIED PURE SAMPLES AS THEY REACT. BECAUSE YOU'RE LOOKING AT REAL PHOTOS OF A REACTION, YOU CAN USE OTHER INFORMATION BEYOND JUST COLOR AND TIMING TO IDENTIFY YOUR SUBSTANCE. DID THE POWDER DISSOLVE QUICKLY OR SLOWLY? WERE THERE LARGE BUBBLES OR SMALL RAPID ONES? ETC...

THE VIDEOS USED TO MAKE OUR CHARTS ARE AVAILABLE ON OUR VIMEO / ANDROID APP / AND BUNKPOLICE.COM





...QUESTION EVERYTHING... @BUNKPOLICE / BUNKPOLICE.COM