


# InertSearch™ for LC

Inertsil® Applications

## Analysis of Ethylenediaminetetraacetic acid ferric sodium salt trihydrate (Under the Condition of food additive analysis method in food, Calcium Disodium Ethylenediaminetetraacetate and Disodium Ethylenediaminetetraacetate)

Data No. LB850-7111



### Conditions

System	: Chromaster HPLC system (HITACHI)
Column	: InertSustain C18 (GL Sciences Inc.) (5 μ m, 250 x 4.6 mm I.D.)
Column Cat. No.	: 5020-07346
Eluent	: A) H <sub>2</sub> O B) CH <sub>3</sub> OH C) Buffer* A/B/C = 78/12/10**, v/v/v
Flow Rate	: 0.8 mL/min
Col. Temp.	: 40 °C
Detection	: UV 254 nm
Injection Vol.	: 20 μ L
Sample	: Standard

### Analyte:

1. Fe(III)-EDTA \*\*\* 10 μ g/mL  
\*\*\*Ethylenediaminetetraacetic acid ferric sodium salt trihydrate

\*Phosphate buffer solution (0.2 mol/L, pH 4.0): Weigh 27.2 g of potassium dihydrogen phosphate, add water to make 1000 mL, and make a 0.2 mol/L potassium dihydrogen phosphate solution. Weigh 23.1 g of phosphoric acid, add water to make 1000 mL, and make a 0.2 mol/L phosphoric acid solution. 0.2 mol/L potassium dihydrogen phosphate solution is added to the 0.2 mol/L phosphoric acid solution to adjust pH to 4.0.

\*\*Weigh 1.61 g of tetra-n-butylammonium bromide, add water/methanol/phosphate buffer (0.2 mol/L, pH 4.0) mixture (78:12:10), and dissolve to make 1000 mL.