

The Posttraumatic Stress Disorder Checklist (PCL)

The PCL (Weathers et al, 1993) is an easily administered self-report rating scale for assessing the 17 DSM-IV symptoms of PTSD. It has excellent test-retest reliability over a 2-3 day period. Internal consistency is very high for each of the three groups of items corresponding to the DSM-IV symptom clusters as well as for the full 17-item scale. The PCL correlates strongly with other measures of PTSD, such as the Mississippi Scale, the PK scale of the MMPI-2, and the Impact of Events Scale, and also correlates moderately with level of combat exposure.

Three versions of the PCL are available, although the differences are very small. The PCL-M is a military version and questions refer to “*a stressful military experience*”. The PCL-S is a non-military version that can be referenced to any specific traumatic event; the questions refer to “the stressful experience”. The PCL-C is a general civilian version that is not linked to a specific event; the questions refer to “a stressful experience from the past”. The scoring is the same for all three versions.

A total score is computed by adding the 17 items, so that possible scores range from 17 to 85. Used as a continuous measure, the PCL has good diagnostic utility. In Vietnam combat veterans a cut-off of 50 on the PCL is a good predictor of a PTSD diagnosis based on the SCID PTSD module. Principal components analysis revealed one large factor, consisting primarily of re-experiencing and hyperarousal items, and one much small factor, consisting primarily of emotional numbing items.

References:

- Blanchard, E.B., Jones-Alexander, J., Buckley, T.C., & Forneris, C.A. (1996). Psychometric properties of the PTSD Checklist (PCL). *Behaviour Research and Therapy*, 34, 669-673.
- Cardove, M.J., Andrykowski, M.A., Redd, W.H., Kenady, D.E., McGrath, P.C., & Sloan, D.A. (1995). Frequency and correlates of posttraumatic stress disorder like symptoms after treatment for breast cancer. *Journal of Consulting and Clinical Psychology*, 63, 981-986.
- Forbes, D., Creamer, M., and Biddle, D. (2001). The validity of the PTSD checklist as a measure of symptomatic change in combat-related PTSD. *Behavior Therapy and Research*, 39, 977-986.
- Weathers, F.W., Litz, B.T., Herman, D.S., Huska, J.A. & Keane, T.M. (1993) The PTSD Checklist (PCL): Reliability, validity, and diagnostic utility. Paper presented at the 9th Annual Conference of the ISTSS, San Antonio.

Common assessment measures: PTSD Checklist

PTSD CheckList – Civilian Version (PCL-C)

Patient's Name: _____

Instructions: Below is a list of problems and complaints that people sometimes have in response to stressful life experiences. Please read each one carefully, put an "X" in the box to indicate how much you have been bothered by that problem *in the past month*.

No.	Response:	Not at all (1)	A little bit (2)	Moderately (3)	Quite a bit (4)	Extremely (5)
1.	Repeated, disturbing <i>memories, thoughts, or images</i> of a stressful experience from the past?					
2.	Repeated, disturbing <i>dreams</i> of a stressful experience from the past?					
3.	Suddenly <i>acting or feeling</i> as if a stressful experience <i>were happening again</i> (as if you were reliving it)?					
4.	Feeling <i>very upset</i> when <i>something reminded</i> you of a stressful experience from the past?					
5.	Having <i>physical reactions</i> (e.g., heart pounding, trouble breathing, or sweating) when <i>something reminded</i> you of a stressful experience from the past?					
6.	Avoid <i>thinking about</i> or <i>talking about</i> a stressful experience from the past or avoid <i>having feelings</i> related to it?					
7.	Avoid <i>activities or situations</i> because <i>they remind you</i> of a stressful experience from the past?					
8.	Trouble <i>remembering important parts</i> of a stressful experience from the past?					
9.	Loss of interest in things that you used to enjoy?					
10.	Feeling <i>distant</i> or <i>cut off</i> from other people?					
11.	Feeling <i>emotionally numb</i> or being unable to have loving feelings for those close to you?					
12.	Feeling as if your <i>future</i> will somehow be <i>cut short</i> ?					
13.	Trouble falling or staying asleep?					
14.	Feeling irritable or having angry outbursts?					
15.	Having difficulty concentrating?					
16.	Being " <i>super alert</i> " or watchful on guard?					
17.	Feeling <i>jumpy</i> or easily startled?					

Weathers, F.W., Huska, J.A., Keane, T.M. *PCL-C for DSM-IV*. Boston: National Center for PTSD – Behavioral Science Division, 1991.

This is a Government document in the public domain.

PTSD CheckList – Civilian Version (PCL-S)

Patient's Name: _____

Instructions: Below is a list of problems and complaints that people sometimes have in response to stressful life experiences. Please read each one carefully, put an "X" in the box to indicate how much you have been bothered by that problem *in the past month*.

The event you experienced was _____ on _____ (date)

No.	Response:	Not at all (1)	A little bit (2)	Moderately (3)	Quite a bit (4)	Extremely (5)
1.	Repeated, disturbing <i>memories, thoughts, or images</i> of a stressful experience from the past?					
2.	Repeated, disturbing <i>dreams</i> of a stressful experience from the past?					
3.	Suddenly <i>acting or feeling</i> as if a stressful experience <i>were happening again</i> (as if you were reliving it)?					
4.	Feeling <i>very upset</i> when <i>something reminded</i> you of a stressful experience from the past?					
5.	Having <i>physical reactions</i> (e.g., heart pounding, trouble breathing, or sweating) when <i>something reminded</i> you of a stressful experience from the past?					
6.	Avoid <i>thinking about</i> or <i>talking about</i> a stressful experience from the past or avoid <i>having feelings</i> related to it?					
7.	Avoid <i>activities or situations</i> because <i>they remind you</i> of a stressful experience from the past?					
8.	Trouble <i>remembering important parts</i> of a stressful experience from the past?					
9.	Loss of interest in things that you used to enjoy?					
10.	Feeling <i>distant</i> or <i>cut off</i> from other people?					
11.	Feeling <i>emotionally numb</i> or being unable to have loving feelings for those close to you?					
12.	Feeling as if your <i>future</i> will somehow be <i>cut short</i> ?					
13.	Trouble falling or staying asleep?					
14.	Feeling irritable or having angry outbursts?					
15.	Having difficulty concentrating?					
16.	Being " <i>super alert</i> " or watchful on guard?					
17.	Feeling <i>jumpy</i> or easily startled?					

Weathers, F.W., Huska, J.A., Keane, T.M. *PCL-S for DSM-IV*. Boston: National Center for PTSD – Behavioral Science Division, 1991.

This is a Government document in the public domain.

PTSD CheckList – Military Version (PCL-M)

Patient's Name: _____

Instructions: Below is a list of problems and complaints that people sometimes have in response to stressful military experiences. Please read each one carefully, put an "X" in the box to indicate how much you have been bothered by that problem *in the past month*.

No.	Response:	Not at all (1)	A little bit (2)	Moderately (3)	Quite a bit (4)	Extremely (5)
1.	Repeated, disturbing <i>memories, thoughts, or images</i> of a stressful experience from the past?					
2.	Repeated, disturbing <i>dreams</i> of a stressful experience from the past?					
3.	Suddenly <i>acting or feeling</i> as if a stressful experience <i>were happening again</i> (as if you were reliving it)?					
4.	Feeling <i>very upset</i> when <i>something reminded</i> you of a stressful experience from the past?					
5.	Having <i>physical reactions</i> (e.g., heart pounding, trouble breathing, or sweating) when <i>something reminded</i> you of a stressful experience from the past?					
6.	Avoid <i>thinking about</i> or <i>talking about</i> a stressful experience from the past or avoid <i>having feelings</i> related to it?					
7.	Avoid <i>activities or situations</i> because <i>they remind you</i> of a stressful experience from the past?					
8.	Trouble <i>remembering important parts</i> of a stressful experience from the past?					
9.	Loss of interest in things that you used to enjoy?					
10.	Feeling <i>distant</i> or <i>cut off</i> from other people?					
11.	Feeling <i>emotionally numb</i> or being unable to have loving feelings for those close to you?					
12.	Feeling as if your <i>future</i> will somehow be <i>cut short</i> ?					
13.	Trouble falling or staying asleep?					
14.	Feeling irritable or having angry outbursts?					
15.	Having difficulty concentrating?					
16.	Being " <i>super alert</i> " or watchful on guard?					
17.	Feeling <i>jumpy</i> or easily startled?					

Weathers, F.W., Huska, J.A., Keane, T.M. *PCL-C for DSM-IV*. Boston: National Center for PTSD – Behavioral Science Division, 1991.

This is a Government document in the public domain.



Using the PTSD Checklist (PCL)

What is the PCL?

The PCL is a 17-item self-report measure reflecting DSM-IV symptoms of PTSD. The PCL has a variety of clinical and research purposes, including:

- screening individuals for PTSD
- aiding in diagnostic assessment of PTSD
- monitoring change in PTSD symptoms

There are three versions of the PCL:

- The **PCL-M** (military) asks about symptoms in response to "stressful military experiences." It is often used with active service members and Veterans.
- The **PCL-C** (civilian) asks about symptoms in relation to generic "stressful experiences" and can be used with any population. This version simplifies assessment based on multiple traumas because symptom endorsements are not attributed to a specific event. In many circumstances it is advisable to also assess traumatic event exposure to ensure that a respondent has experienced at least one event that meets DSM-IV Criterion A.
- The **PCL-S** (specific) asks about symptoms in relation to an identified "stressful experience." The PCL-S aims to link symptom endorsements to a specified event. Similar to the PCL-C, it is optimal to assess traumatic event exposure to ensure that the index event meets PTSD Criterion A. Respondents also can be instructed to complete the PCL-S in reference to a specified event or event type (e.g., assault, disaster, or accident).

How is the PCL administered and scored?

The PCL is a self-report instrument that can be read by respondents themselves or read to them either in person or over the telephone. It can be completed in approximately 5-10 minutes.

The PCL can be scored in several ways:

- A **total symptom severity score** (range = 17-85) can be obtained by summing the scores from each of the 17 items that have response options ranging from 1 "Not at all" to 5 "Extremely".
- The gold standard for diagnosing PTSD is a structured clinical interview such as the Clinician-Administered PTSD Scale (CAPS). When necessary, the PCL can be scored to provide a presumptive diagnosis. This has been done in three ways:
 - (1) determine whether an individual meets DSM-IV symptom criteria as defined by at least 1 B item (questions 1-5), 3 C items (questions 6-12), and at least 2 D items (questions 13-17).

Symptoms rated as “Moderately” or above (responses 3 through 5 on individual items) are counted as present.

- (2) determine whether the total severity score exceeds a given normative threshold (see Table 1).
- (3) combine methods (1) and (2) to ensure that an individual meets both the symptom pattern and severity threshold.

Choosing a cut-point score

Factors to be considered when choosing a PCL cut-point score include:

- **The goal of the assessment:** A lower cut-point is considered when screening for PTSD or when it is desirable to maximize detection of possible cases. A higher cut-point is considered when informing diagnosis or to minimize false positives.
- **The prevalence of PTSD in the target setting:** Generally, the lower the prevalence of PTSD in a given setting, the lower the optimal cut-point. In settings with expected high rates of PTSD, such as specialty mental health clinics, consider a higher cut-point. In settings with expected low rates of PTSD, such as primary care clinics or circumstances in which patients are reluctant to disclose mental health problems, consider a lower cut-point.

Below are suggested cut-point ranges based on prevalence and setting characteristics. There is no absolute method for determining the correct cut-point on the PCL. If you know the prevalence of PTSD in your target population, use column 1 to find the suggested PCL cut-point (column 3). If you do not know the prevalence in your population, you can choose a cut-point based on the type of setting (column 2) in which you are working. Consider scores on the low end of the range if the goal is to screen for PTSD. Consider scores on the high end of the range if the goal is to aid in diagnosis of PTSD.

Table 1. Suggested PCL cut-point scores

Estimated Prevalence of PTSD	Typical Setting	Suggested PCL Cut-Point Scores
15% or Below	e.g. civilian primary care, Department of Defense screening, or general population samples	30-35
16-39%	e.g. specialized medical clinics (such as TBI or pain) or VA primary care	36-44
40% or Above	e.g. VA or civilian specialty mental health clinics	45-50

Note. These recommendations are general and approximate, and are not intended to be used for legal or policy purposes. Research is needed to establish optimal cut-point scores for a specific application.

Measuring change

Good clinical practice often involves monitoring patient progress. Evidence suggests that a 5-10 point change is reliable (i.e., not due to chance) and a 10-20 point change is clinically meaningful (Monson et al., 2008). Therefore, we recommend using 5 points as a minimum threshold for determining whether an individual has responded to treatment and 10 points as a minimum threshold for determining whether the improvement is clinically meaningful.

Studies that informed our recommendations

1. Blanchard, E. B., Jones-Alexander, J., Buckley, T. C., & Forneris, C. A. (1996). Psychometric properties of the PTSD checklist (PCL). *Behavioral Research & Therapy*, 34, 669-673.
2. Bliese, P. D., Wright, K. M., Adler, A. B., Cabrera, O., Castrol, C. A., & Hoge, C. W. (2008). Validating the primary care posttraumatic stress disorder screen and the posttraumatic stress disorder checklist with soldiers returning from combat. *Journal of Consulting and Clinical Psychology*, 76, 272-281.
3. Dobie, D. J., Kivlahan, D. R., Maynard, C., Bush, K. R., McFall, M. E., Epler, A. J., & Bradley, K. A. (2002). Screening for post-traumatic stress disorder in female veteran's affairs patients: Validation of the PTSD checklist. *General Hospital Psychiatry*, 24(6), 367-374.
4. Freedy, J. R., Steenkamp, M. M., Magruder, K. M., Yeager, D. E., Zoller, J. S., Hueston, W. J., & Carek, P. J. (2010). Post-traumatic stress disorder screening test performance in civilian primary care. *Family Practice*.
5. Harrington, T., & Newman, E. (2007). The psychometric utility of two self-report measures of PTSD among women substance users. *Addictive Behaviors*, 32, 2788-2798.
6. Keen, S. M., Kutter, C. J., Niles, B. L., & Krinsley, K. E. (2008). Psychometric properties of PTSD checklist in sample of male veterans. *Journal of Rehabilitation Research and Development*, 45(3), 465-474.
7. Monson, C. M., Gradus, J. L., Young-Xu, Y., Schnurr, P. P., Price, J. L., & Schumm, J. A. (2008). Change in posttraumatic stress disorder symptoms: Do clinicians and patients agree? *Psychological Assessment*, 20(2), 131-138.
8. Kimerling, R., Prins, A., Yeager, D.E., & Magruder, K.M. (2010, November). An interval approach to screening for PTSD in primary care. Poster presented at the 44th annual conference of the Association for Cognitive and Behavior Therapy, San Francisco, CA.
9. Sherman, J. J., Carlson, C., Wilson, J. F., Okeson, J., & McCubbin, J. A. Posttraumatic stress disorder among patients with orofacial pain. *Journal of Orofacial Pain*, 19, 309-317.
10. Walker, E. A., Newman, E., Dobie, D. J., Ciechanowski, P., & Katon, W. (2002). Validation of the PTSD checklist in an HMO sample of women. *General Hospital Psychiatry*, 24, 375-380.
11. Weathers, F., Litz, B., Herman, D., Huska, J., & Keane, T. (October 1993). *The PTSD Checklist (PCL): Reliability, Validity, and Diagnostic Utility*. Paper presented at the Annual Convention of the International Society for Traumatic Stress Studies, San Antonio, TX. NOTE: Due to some confusion over versions of the PCL, some of the published papers state that the PCL-C was used in this study, but the authors have confirmed that the PCL-S was the version actually used.
12. Yeager, D. E., Magruder, K. M., Knapp, R. G., Nicholas, J. S., & Frueh, B. C. (2007). Performance characteristics of the posttraumatic stress disorder checklist and SPAN in Veterans affairs primary care settings. *General Hospital Psychiatry*, 29(4), 294-301.