


# InertSearch for LC

Inertsil Applications

## Analysis of Paracetamol and Mefenamic acid

(Under the Condition of the draft for the Indian Pharmacopoeia, Paracetamol and Mefenamic Acid Suspension)

Data No. LB882-7111



### Conditions

**System** : Chromaster HPLC system (HITACHI)  
**Column** : InertSustain C18 (GL Sciences Inc.)  
(5  $\mu$  m, 250 x 4.6 mm I.D.)  
**Column Cat. No.** : 5020-07346  
**Eluent** : A) CH<sub>3</sub>CN  
B) CH<sub>3</sub>OH  
C) Buffer\*  
A/B/C = 40/10/50, v/v/v  
**Flow Rate** : 1.0 mL/min  
**Col. Temp.** : 25 °C  
**Detection** : UV 285 nm  
**Injection Vol.** : 10  $\mu$  L  
**Sample** : Standard

### Analyte:

1. Paracetamol	0.125 mg/mL
2. Mefenamic acid	0.05 mg/mL
Tailing factor	
peak area of 1	: 1.16 ( $\leq$ 2.0)
peak area of 2	: 1.20 ( $\leq$ 2.0)
RSD of the	
peak area of 1 (%)(n=6)	: 0.27 ( $\leq$ 2.0)
peak area of 2 (%)(n=6)	: 0.32 ( $\leq$ 2.0)

\* Dissolve 8.37g of potassium dihydrogen orthophosphate and 6.71g of dipotassium hydrogen orthophosphate in 1000 ml of water, adjusted to pH 6.5 with dilute orthophosphoric acid.