

We will introduce an example of the antioxidant capacity measurement of food using post-column HPLC using 1,1-diphenyl-2-picrylhydrazyl free radical (DPPH), which is known as a stable radical.


DPPH is known as a purple stable radical and becomes colorless when reduced by antioxidants.

DPPH radical scavenging activity is known as an index of antioxidant substances that utilize this reaction. By measuring this, it is possible to relatively evaluate the antioxidant capacity. This time, we performed measurements together with UV direct detection, so we will introduce them. There is a known method of calculating by converting to the amount of Trolox, but we will omit it here.

(K. Suzuki)

### Example of standard solution analysis

It can be confirmed that catechin has DPPH activity. On the other hand, we can see that the caffeine peak has very low DPPH activity despite having a higher UV absorption.



#### HPLC Condition

**Column** : InertSustain C18  
(5  $\mu$ m, 150  $\times$  4.6 mm I.D.)

**Cat. No.** : 5020-07345

**Eluent** : A) 0.1%  $\text{H}_3\text{PO}_4$  in  $\text{H}_2\text{O}^*$   
B)  $\text{CH}_3\text{OH}/\text{CH}_3\text{CN} = 9/1$ , v/v

Time(min)	A(vol%)	B(vol%)
0.0	90	10
15.0	80	20
30.0	60	40
30.1	90	10
40.0	90	10

\*1 mL of special grade phosphoric acid (85%) added to 1 L of radical ultrapure water

**Flow rate** : 1.0 mL/min

**Col.Temp.** : 40  $^\circ\text{C}$

**Detection** : UV 280 nm (UV)  
VIS 520 nm (DAD)


**Inj.Vol.** : 10  $\mu\text{L}$

**Reaction solution** : 50 mg/L DPPH radical in MeOH

**Reaction flow rate** : 0.5 mL/min

**Reaction temperature** : 40  $^\circ\text{C}$

#### Structure




Structures are created using Chemistry 4-D Draw which is provided by ChemInnovation Software, Inc.


## Analysis example

Using LC Technical Note No. 145 as a reference, two types of tea were passed through a 0.45  $\mu\text{m}$  GL chromatodisk and measured.


### Green Tea A



### Green Tea B



## Flow diagram



GL Sciences disclaims any and all responsibility for any injury or damage which may be caused by this data directly or indirectly. We reserve the right to amend this information or data at any time and without any prior announcement.

### GL Sciences Inc. Japan

22-1 Nishishinjuku 6-chome  
 Shinjuku-ku, Tokyo  
 163-1130, Japan

Phone: +81-3-5323-6620  
 Fax: +81-3-5323-6621  
 Email: [world@glsc.co.jp](mailto:world@glsc.co.jp)  
 Web: [www.glsciences.com](http://www.glsciences.com)

### GL Sciences Inc. USA

4733 Torrance Blvd. Suite 255  
 Torrance, CA 90503  
 USA

Phone: +1-310-265-4424  
 Fax: +1-310-265-4425  
 Email: [info@glsciencesinc.com](mailto:info@glsciencesinc.com)  
 Web: [www.glsciencesinc.com](http://www.glsciencesinc.com)

### GL Sciences B.V.

Dillenburgstraat 7C  
 5652AM, Eindhoven  
 The Netherlands

Phone: +31-40-254-9531  
 Email: [info@glsciences.eu](mailto:info@glsciences.eu)  
 Web: [www.glsciences.eu](http://www.glsciences.eu)

### GL Sciences (Shanghai) Limited

Tower B, Room 2003  
 Far East International Plaza  
 No.317 Xianxia Road, Changning District  
 Shanghai, China 200051

Phone: +86-21-62782272  
 Email: [contact@glsciences.com.cn](mailto:contact@glsciences.com.cn)  
 Web: [www.glsciences.com.cn](http://www.glsciences.com.cn)



International Distributors

Visit our Website at [www.glsciences.com/distributors](http://www.glsciences.com/distributors)